WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5:

C07H 19/073, A61K 31/70

(11) International Publication Number:

WO 92/08727

A1

(43) International Publication Date:

29 May 1992 (29.05.92)

(21) International Application Number:

PCT/EP91/02134

(22) International Filing Date:

9 November 1991 (09.11.91)

(30) Priority data:

22032 A/90

13 November 1990 (13.11.90) IT

(71) Applicants (for all designated States except US): CONSIGLIO NAZIONALE DELLE RICERCHE [IT/IT]; Piazzale Aldo Moro, 7, 1-00195 Rome (IT). MENARINI RICERCHE SUD S.P.A. [IT/IT]; Via Tito Speri, 10, I-0040 Bernaria (IT). 00040 Pomezia (IT).

(72) Inventors; and

(72) Inventors; and
(75) Inventors/Applicants (for US only): IOTTI, Stefano [IT/IT];
Via E. Torricelli, 23, I-42100 Reggio Emilia (IT). CO-LONNA, Francesco, Paolo [IT/IT]; Via San Felice, 161,
I-40122 Bologna (IT). GARBESI, Anna, Maria [IT/IT];
Via Bonavia, 12, I-40068 San Lazzaro Savena (IT).
SPADARI, Silvio [IT/IT]; Via Calabria, 6, I-27100 Pavia
(IT). FOCHER, Federico [IT/IT]; Via Don Minzoni, 5,
I-27100 Pavia (IT). CIARROCCHI, Giovanni [IT/IT];
Piazza Castello, 8, I-27040 Cigognola (IT). ARCA-MONE, Federico [IT/IT]; Via 4 Novembre, 26, I-20014
Nerviano (IT). Nerviano (IT).

(74) Agent: GERVASI, Gemma; Notarbartolo & Gervasi S.r.l., Viale Bianca Maria, 33, I-20122 Milan (IT).

(81) Designated States: AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH (European patent), CI (OAPI patent), CM (OAPI patent), CS, DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GA (OAPI patent), GB (European patent), GN (OAPI patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU (European patent), MC, MG, ML (OAPI patent). MN. MR (OAPI patent). MW. NL ML (OAPI patent), MN, MR (OAPI patent), MV, MU, (European patent), NO, PL, RO, SD, SE (European patent), SN (OAPI patent), SU+,TD (OAPI patent), TG (OAPI patent), US.

Published

With international search report.

(54) Title: L-2'-DESOXYURIDINES AND PHARMACEUTICAL COMPOSITIONS CONTAINING THEM

(57) Abstract

4

L-2'-desoxyuridines of formula (I) are described, wherein R=H, CH₂OC₂H₅, CH₂OH, CH₃, -CH=CBrH and R'=OH, F, N₃, and their use for the preparation of pharmaceutical compositions useful for the treatment of viral infections in men and in animals.

+ DESIGNATIONS OF "SU"

Any designation of "SU" has effect in the Russian Federation. It is not yet known whether any such designation has effect in other States of the former Soviet Union.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria-	ES	Spain	MG	Madagascar
UA	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	CB	United Kingdom	MW	
BG	Bulgaria	GN	Guinea		Malawi
BJ	Benin	GR:	Greece	NL	Netherlands
BR	Brazil	HU	Hungary	NO	Norway
CA	Canada .	17	ltaly	. PL	Poland
CF	Central African Republic	JP	-	RO	Romania
CC	Congo	KP	Japan	SD	Sudan
CH	Switzerland	KP	Democratic People's Republic	SE	Sweden
CI.		••-	of Korea	SN	Senegal
	Côte d'Ivoire	KR	Republic of Korea	ຣບ+	Soviet Union
CM	Cameroon	LI	Licchtenstein	TD	Chad
cs	Czuchoslovakia	LK	Sri Lanka	TC	Togo
DE	Germany	LU.	Luxembourg	US	United States of America
DK	Denmark	MC	Monaco		

L-2'-DESOXYURIDINES AND PHARMACEUTICAL COMPOSITIONS CONTAINING THEM.

Invention field

The present invention relates to L-2'-desoxyuridine of general formula (I)

10

wherein R = H, $CH_2OC_2H_5$, CH_2OH , CH_3 , -CH=CBrH and R' = OH, F, 15 N_3 and their use for the preparation of pharmaceutical compositions having antiviral properties.

Prior Art

L-2'-desoxynucleosides of formula (I) in which R = H, CH₃, CH₂OH and R' = OH are known and their synthesis was effected as described in the literature [Coll. Czech. Chem. Comm. <u>37</u>, p. 4072 (1972)] starting from L-arabinose and cyanamid as shown hereinbelow, R and R' being as defined supra, H excluded:

It is known furthermore that derivatives of 2'-desoxyuridine belonging to the natural series D are phosphorylated by viral thymidino-kinase [Y.C. Cheng, "Antimetabolites in Biochemistry, Biology and Meidine", Pergamon Press (1979)]. In "Nucleic Acid Research" 1976, 3 (8), 2143-54 (Eng) is reported that L-uridine, L-cytidine and L-thymidine when administered to mice are largely eliminated in the urine in unmodified form leaving in some tissues only traces of phosphorylated metabolites.

In the light of the cited literature, therefore, it was neither known nor could be foreseen that also non-natural 2'-desoxyuridines of the L-series would behave similarly to the natural compounds of the D-series, and even less that they could be employed in the pharmaceutical field.

15 Detailed description of the invention

We have now surprisingly found, and this is an object of the present invention, that compounds of general formula (I) in which R = H, $CH_2OC_2H_5$, CH_2OH , CH_3 , -CH=CBrH and R' = OH, F, N_3 , are substrates of HSV 1 and HSV 2 thymidino-kinase (that is they are phosphorylated by the viral enzyme) while they are wholly inert to the action of mammalian thymidino-kinase.

The phosphorylation of these compounds by the viral enzyme is of considerable importance, allowing to obtain drugs capable of inhibiting the development of pathogenous viruses. In fact, in the infected cell, the analogous monophosphates thus formed

undergo a successive transformation into di-phosphates by the action of the viral thymidino-kinase (TK) and then to triphosphate nucleotides by action of unspecific cellular enzymes. Such analogous triphosphates behave as selective inhibitors of viral DNA polymerases, or, more frequently, as improved substrates, being preferentially incorporated in the viral DNA destroying its functionality.

In the healthy cell, where, the activation of the nonnatural analogous compounds to mono- and di-phosphates does not take place, neither does their transformation to triphosphates, with a consequent sharp reduction of the toxic effects in the areas not affected by the viral infection.

Particularly the present invention relates to pharmaceutical compositions having antiviral properties containing as active principle an L-2'-desoxyuridine of general formula (I), wherein R = H, CH₂OC₂H₅, CH₂OH, CH₃, -CH=CBrH and R' = OH, F, N₃ and a pharmaceutically acceptable excipient or vehicle. The compounds of formula (I) wherein R = -CH=CBrH, CH₃, and R' = OH, F, N₃, with the proviso that when R = CH₃, R' is not OH, are new.

According to the present invention, the preferred compounds of general formula (I) suitable for the above said purposes are:

- 1-(2-desoxy-β-L-erythro-pentafuranosyl)-2,4-(1H,3H)

 pyrimidindione (R = H, R' = OH)
- 25 1-(2-desoxy- β -L-erythro-pentafuranosyl)-5-hydroxymethylen-2,4-(1H,3H)-pyrimidindione (R = CH₂OH, R' = OH)

- 1-(2-desoxy- β -L-erythro-pentafuranosyl)-5-ethoxymethylen-2,4-(1H,3H)-pyrimidindione (R = CH₂OC₂H₅, R' = OH)
- 1-(2-desoxy- β -L-erithro-pentafuranosyl)-5-methyl-2,4-(1H,3H)-pyrimidindione (R = CH₂, R' = OH)
- 5 1-(2-desoxy-β-L-erythro-pentafuranosyl)-5-(E-2-bromovinyl)2,4-(1H,3H)-pyrimidindione (R = -CH=CBrH, R' = OH)
 - 1-(2,3-didesoxy-3-azido- β -L-erythro-pentafuranosyl)-5-methyl-2,4-(1H,3H)-pyrimidindione (R = CH₃, R' = N₃)
 - 1-(2,3-didesoxy-3-fluoro-β-L-erythro-pentafuranosyl)-5-
- methyl-2,4-(1H,3H)-pyrimidindione (R = CH₃, R' = F)

 The first four compounds were described in the literature without, however, any reference to their antiviral properties and their ensuing pharmaceutical use, while the others are new. For illustrative, non limiting, purposes, we are now reporting the preparation of one compound according to the present invention.

EXAMPLE 1

Synthesis of 1-(2-desoxy- β -L-erythro-pentafuranosyl)-5-methyl-2,4(1H,3H)-pyrimidindione (R = CH₃, R' = OH).

- 20 A mixture of 4.6 g L-2'-desoxyuridine (compound of formula (I) in which R = H, R' = OH), 20 ml 1 M KOH and 20 ml aqueous 37% formaldehyde is kept at 60°C for five days, adding every 24 hours 5 ml 1 M KOH and 5 ml aqueous 37% formaldehyde. The reaction mixture is then diluted with an equal volume of water,
- 25 the pH is brought to 3 by addition of Dowex 50 (H+) ion

exchange resin, then filtered washing the resin with 200 ml water; filtrate and washing water are put together and evaporated under reduced pressure. The residue is then coevaporated for three times with absolute ethanol, dissolved in 50 ml of same solvent, made alkaline by addition of triethylamine, evaporated under reduced pressure and dried by co-evaporation with 50 ml toluene (twice). The residue is kept for one night under reduced pressure over phosphoric anhydride, then 200 ml absolute ethanol are added and the solution obtained is brought to pH 2.5 by addition of concentrated hydrochloric acid and heated on reflux for 2 hrs.

After cooling, the reaction mixture is made alkaline by addition of triethylamine and evaporated; the residue is chromatographed on silica gel eluting with a methylene chloride/methanol 85ml/15ml mixture.

The obtained compound is then dissolved in 200 ml absolute ethanol, 0.5 ml concentrated hydrochloric acid and 1 g palladium 10% on carbon are added, followed by hydrogenation at normal pressure for 3 hrs. The reaction mixture is filtered on 20 Celite, which is washed with 100 ml ethanol. The filtrate are put together, made alkaline with triethylamine and evaporated under reduced pressure. The residue is crystallized from 8 ml absolute ethanol.

Thus 2 g L-thymidine $[\alpha]$ = -18.5° (c = 1% in water).

25 With a method similar to the one described in the literature for the compounds derived from D-2'-desoxyuridine, and

employing the suitable reagents, one can obtain also the other previously indicated compounds of formula (I).

Biological activity

The biolgical activity of the compounds according to the present invention in the treatment of viral infections was evaluated employing HSV 1 TK and cell TK.

Fig. 1 shows the L-thymidine effect on the enzymathic activities of Herpes simplex 1 virus TK and of the human one.

Fig. 1 shows that L-thymidine inhibits the phosphorylation of 10 (D)-thymidine by Herpes simplex 1 TK (\square), but not by human TK (\spadesuit).

The test consists in measuring the amount of (D)-thymidine substrate phosphorylated to (D)-TMP (ordinates) in the presence of growing L-thymidine concentrations (abscissae). The viral or cellular purified enzyme (0.07 units) was incubated for 15 minutes at 37°C in 25 µl of a mixture containing 30 mM Hepes-K, pH 7.5, 6 mM MgCl₂, 6 mM ATP, 0.5 mM dithiothreitol (DTT), 10 µM [³H]Thy (25 Ci/mmole) and various L-thymidine concentrations.

20 The reaction is stopped depositing 20 µl of the mixture on DE-81 filters which are immediately soaked in an excess of 1 mM ammonium formiate, pH 5.6, in order to eliminate the (D)-[3H] thymidine not transformed into monophosphate and which cannot therefore bind itselft with the positive charges of the DE-81 filter.

The filters are then washed in distilled water for 5 minutes and dehydrated in ethyl alcohol for 5 minutes. The radioactive D-TMP bound to the filter was evaluated in a β radiation counter.

- One can clearly observe a specific inhibition of viral TK; in fact at 5 μ g/ml viral TK is 90% inhibited, whereas the human one is completely resistent. Human TK proved to be fully resistent to L-thymidine even at the highest investigated dose (200 μ g/ml).
- L-thymidine competes with D-thymidine for the active site of the viral enzyme, as proved by the competitive type inhibition curves reported in Fig. 2, which is a representation according to Lineweaver-Burk of the effect of L-thymidine on the TK activity of Herpes simplex virus in the presence of different concentrations of D-[3H] thymidine substrate (expressed in the abscissae as the inverse of concentration).

The viral enzyme (0.07 units) was incubated for 15 minutes at 37°C in 25 µl of a mixture containing 30 mM Hepes-K, pH 7.5, 6 mM MgCl₂, 6 mM ATP, 0.5 mM dithiothreitol (DTT), various concentrations of D-[³H]thymidine(25 Ci/mmol) and various concentrations of L-thimidine (0 µM [5], 2 µM [4], 5 µM [7], 10 µM [4], 15 mM [7]). The reaction is stopped by depositing 20 µl of the mixture on DE-81 filters which are processed as indicated in Fig. 1. The D-TMP values are reported as the inverse of the concentration on the ordinates. From the experiment one concludes that Km for D-thymidine viral enzyme

is 2.8 µM and that Ki for L-thymidine is 2 µM.

To check whether L-thymidine is phosphorylated by the viral enzyme, as happens for D-thymidine, IDU, TFT, ACU etc., or whether it simply inhibits the synthesis of the natural substrate competing for the active site, as it happens for the other compounds which we have studied, such as phenyldesoxyguanosine (Ph-dG) we have analysed by HPLC the products of the reaction catalyzed by viral TK, between $[\gamma^{-32}P]$ ATP and L-thymidine.

The investigated nucleosides and nucleotides were separated by the reverse-phase method employing the Bio-Rad 100 MAPS preparative system.

A reverse-phase C₁₈ Bio Sil ODS-5S (0.4 x 15 cm) column was employed under the following conditions: injected volume 20 µl; 15 UV : 260 nm; temperature : room; eluent : buffer A (20 mM KH₂PO₄, pH 5.6), buffer B (20 mM KH₂PO₄, pH 5.6, 60% methanol). The specific conditions for separating L-thiomidine ATP are as follows: from 0 to 20 min. a 0% to 70% buffer B gradient; from 20 to 30 min. a 70% to 77% buffer B gradient and from 30 to 32 min. a 77% to 100% buffer B gradient. Flux was 0.5 ml/min. The enzymatic reaction (0.3 units viral enzyme) was performed as previously described, except that, instead of 6 mM ATP, 100 µM [y -32p] ATP 1500 cpm/pmol was employed and that the 37°C incubation was protracted to 30 min. The results obtained are reported in Fig. 3, in which on the abscissae of each panel A,

20

B and C the number of fractions eluted by the column is reported. In each fraction, the radioactivity was determined, which is reported as counts per minute (cpm) on the respective ordinates.

5 Panel A shows the control data without Thy, panel B the data for 10 μ M L-thymidine and panel C the data for 10 μ M D-thymidine in the assay.

As can be seen from Fig. 3, viral TK phosphorylizes L-thymidine: in fact, in the presence of L-thymidine a radioactivity peak in the same position as D-TMP (Panel C) is obtained.

In the 30 min. reaction, viral TK phosphorylizes 70% of the L-thymidine present in the assay, which is comparable to what is obtained with natural D-thymidine.

15 Pharmaceutical Compositions

The pharmaceutical compositions according to the present invention comprise, as an active component, a therapeutically effective amount of a L-desoxyuridine of general formula (I) in which R and R' have the previously indicated meanings, or one of its pharmaceutically acceptable salts, in association with one or more pharmaceutically acceptable excipients or vehicles. The pharmaceutical compositions according to the invention may be administered per os, parenterally and topically in suitable pharmaceutical formulations, for instance as sterile solutions for injectable use, tablets, capsules, powders, granulates, syrups, colloyria, ointments, creams, suppositories, ovules,

bougies, etc.

The active principle is contained in the pharmaceutical compositions according to the invention in amounts variable between 50 mg and 2 g per dose, depending on the way of administration.

Useful excipients in the formulations according to the invention are e.g. jellying agents, auxiliaries for gelatine capsules, antioxidants, dispersing agents, emulsifiers, antifoam agents, taste correcting agents, preservers, solubilyzing agents, etc.

CLAIMS

1 1. L-2'-desoxyuridines of general formula (I)

- 2 wherein R = -CH = CBrH, CH_3 and R' = OH, $F_1 = N_3$, with the proviso
- 3 that when $R = CH_3$, R' is not OH, and their pharmaceutically
- 4 acceptable salts.
- 1 2. Compound of general formula (I) according to claim 1,
- 2 wherein R = -CH=CBrH and R' = OH and its pharmaceutically
- 3 acceptable salts.
- $_1$ 3. Compound of general formula (I) according to claim 1,
- 2 wherein $R = CH_3$ and $R' = N_3$ and its pharmaceutically acceptable
- g salts.
- 1 4. Compound of general formula (I) according to claim 1,
- 2 wherein R = CH₂ and R' = F and its pharmaceutically acceptable
- 3 salts.
- 1 5. Pharmaceutical compositions containing as active principle
- 2 an effective amount of a compound according to claim 1 and
- 3 pharmaceutically acceptable excipients or vehicles.
- 1 6.Pharmaceutical composition according to claim 5 wherein the
- 2 active principle is a compound of formula (I) wherein R =
- 3 -CH=CBrH and R' = OH and its pharmaceutically acceptable salts.

- 1 7. Pharmaceutical composition according to claim 5 wherein the
- 2 active principle is a compound of formula (I) wherein $R = CH_2$
- 3 and R' = N_3 and its pharmaceutically acceptable salts.
- 1 8. Pharmaceutical composition according to claim 5 wherein the
- 2 active principle is a compound of formula (I) wherein $R = CH_3$
- 3 and R' = F and its pharmaceutically acceptable salts.
- 1 9. Pharmaceutical compositions according to claim 5 having
- 2 antiviral acivity.
- 1 10. Pharmaceutical compositions according to claim 5 for the
- 2 therapeutic treatment of Herpes simplex.
- 1 11. Pharmaceuticals compositions containing as active
- 2 principle an effective amount of a compound of formula (I)

$$\begin{array}{c} O \\ H-N \\ N \\ O \\ \end{array} \begin{array}{c} R \\ \end{array}$$
 OH

- 3 wherein R = H, $CH_2OC_2H_5$, CH_2OH and R' = OH, F, N_3 or R = CH_3
- 4 and R' = OH and their pharmaceutically acceptable salts and
- 5 pharmaceutically acceptable excipients or vehicles.
- 1 12. Pharmaceutical composition according to claim 11 wherein
- 2 the active principle is a compound of formula (I)
- 3 wherein R = H and R' = OH and its pharmaceutically, acceptable
- 4 salts.
- 1 13. Pharmaceutical composition according to claim 11 wherein

- 2 the active principle is a compound of formula (I) wherein R =
- 3 CH2OH and R' = OH and its pharmaceutically acceptable salts.
- 1 14. Pharmaceutical composition according to claim 11 wherein
- 2 the active principle is a compound of formula (I)
- 3 wherein $R = CH_2OC_2H_5$ and R' = OH and its pharmaceutically
- 4 acceptable salts.
- 1 15. Pharmaceutical composition according to claim 11 wherein
- 2 the active principle is a compound of formula (I) wherein R =
- 3 CH_2 and R' = OH and its pharmaceutically acceptable salts.
- 1 16. Pharmaceutical compositions according to claim 11 having
- 2 antiviral activity.
- 1 17. Pharmaceutical compositions according to claim 11 for the
- 2 therapeutic treatment of Herpes simplex
- 1 18. Pharmaceutical compositions according to claims 5 or 11
- 2 wherein the active principle is contained in an amount of
- 3 between 50 mg and 2 g.
- 1 19. Pharmaceutical compositions according to claims 5 or 11.
- 2 for parenteral, oral or topical administration.
- 1 20. Pharmaceutical compositions according to claims 5 or 11 in
- 2 the form of injectable sterile solutions, tablets, capsules.
- 3 granulates, powders, syrups, collyria, ointments, creams,
- 4 suppositories, ovules, bougies.





FIG. 3



International Application No

I. CLASSIF	ICATION OF SUBJE	CCT MATTER (if several classification syn	nbols apply, indicate all) ⁶		
According	o International Patent	Classification (IPC) or to both National Cla	ssification and IPC		
Int.C1.	5 CO7H19/0	73; A61K31/70			
		·			
II. FIELDS	SEARCHED	Minimum Documen	teston Serrehad?		
			Tassification Symbols		
Classificati	on System		iassin carron by in a carron		
Int.Cl.	5	CO7H ; A61K			
		Documentation Searched other to to the Extent that such Documents at	han Minimum Documentation re included in the Fields Searched ⁸	,	
III. DOCU	MENTS CONSIDER	ED TO BE RELEVANT?			
Category °	Citation of D	ocument, 11 with indication, where appropria	te, of the relevant passages 12	Relevant to Claim No.13	
X	1990	352 248 (MEDIVIR AKTIEBO	OLAG) 24 January	1,3-5, 7-12, 15-20	
Υ	FR. A. 2	whole document 627 492 (INSTITUT DE RECOGIQUES APPLIQUEES) 25	CHERCHES CHIMIQUES August 1989	1,3-5, 7-12, 15-20	
	see abs	tract		15-20	
Υ	WO,A,8) 27 Ju	906 655 (UNIVERSITE PIE ly 1989	RRE ET MARIE CURIE	1,3-5, 7-12, 15-20	
	see abs			1 2	
Y	EP,A,O April 1	217 580 (THE WELLCOME F 987	OUNDATION LID.) 8	1,3-5, 7-12, 15-20	
	see cla	ims 4,11	-/		
Special categories of cited documents: 10 **A document defining the general state of the art which is not considered to be of particular relevance **E** earlier document but published on or after the international filing date **It** document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) **O** document referring to an oral disclosure, use, exhibition or other means **P** document published prior to the international filing date but later than the priority date claimed **It** It ater document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention **Coument of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. **E** document member of the same patent family **It** document member of the same patent family					
:[. 16 JAN	WARY 1992	1 5. U.	JC	
Internation	al Searching Authority	ZAN PATENT OFFICE	Signature of Authorized Officer SCOTT J.R. 4.R	M.S. 26171	

Category °	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
		RESERVED TO CIETAL TIO
l		·
Y	EP,A,O 254 268 (AKADEMIE DER WISSENSCHAFTEN DER DDR) 27 January 1988	1,3-5, 7-12, 15-20
.	see abstract	10 20
4	COLLECTION OF CZECHOSLOVAK CHEMICAL COMMUNICATIONS. vol. 37, no. 12, December 1972, PRAGUE CS pages 4072 - 4087; A.HOLY: 'Nucleic Acid Components and their	1,3-5, 7-12, 15-20
	Analogues. CLIII. Preparation of 2'-deoxy - L - Ribonucleosides of the Pyrimidine Series.' cited in the application see abstract see page 4073, compound VII see page 4075, compounds XX, XXI, XXII	
Х,Р	JOURNAL OF ORGANIC CHEMISTRY. vol. 56, no. 11, 24 May 1991, EASTON US pages 3591 - 3594; J.WENGEL ET AL.: 'Synthesis of L-3'-Azido-3'-deoxythymidine and its	1,3,5,7, 9-11,16, 18-20
	Stereoisomers' see the whole document	
-		
		

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO. EP SA 9102134

52789

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 16/01/92

cited in search report	Publication date		Patent family member(s)	Publication date
EP-A-0352248	24-01-90	AU-A- WO-A-	3978689 9001036	19-02-90 08-02-90
FR-A-2627492	25-08 - 89	AU-A-	3213789	22-09-89
		EP-A-	0448550	02-10-91
	4	WO-A-	8908115	08-09-89
		JP-T-	3504492	03-10-91
WO-A-8906655	27-07-89	FR-A-	2626001	21-07-89
,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		FR-A-	2640267	15-06-90
		EP-A-	0325537	26-07-89
•		JP-T-	2503002	20-09-90
EP-A-0217580	08-04-87	AU-B-	595832	12-04-90
		AU-A-	6270286	19-03-87
		GB-A-	2181128	15-04-87
		AT-B-	390000	26-02-90
		AT-B-	392794	10-06-91
		AU-B-	572019	28-04-88
		CA-A-	1238277	21-06-88
		DE-A-	3608606	18-09-86
		DE-A-	3645058	02-02-89
	•	DE-A-	3645059	05-01-89
		EP-A,B	0196185	01-10-86
		EP-A-	0291633	23-11-88
	,	EP-A-	0306597	15-03-89
		JP-A- JP-A-	63290895 62103100	28-11-88 13-05-87
		US-A-	4847244	13-05-87
		US-A-	4847244 4818750	04-04-89
·		US-A-	4857511	15-08-89
	27-01-88	JP-A-	63107936	12-05-88
EF-M-0234200	~1-01-00	US-A-	4963662	16-10-90
		AU-B-	615431	03-10-91
		AU-A-	1259288	10-11-88

FORM POST9

. 1

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82